

## University of Groningen

### Caprolactam from renewable resources

Teddy, T

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2013

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Teddy, T. (2013). *Caprolactam from renewable resources: catalytic conversion of 5-hydroxymethylfurfural into caprolactone*. s.n.

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Stellingen

Behorende bij het proefschrift:

## Caprolactam from Renewable Resources: Catalytic Conversion of 5-Hydroxymethylfurfural into Caprolactone

Teddy

1. The use of extremely high pressures (380 bar) for the reaction of 5-hydroxymethylfurfural (HMF) to 1,6-hexanediol is not necessary.

*T. Utne, J. D. Garber, R. E. Jones, US Patent 3083236, 1963.*

2. Yield data without information on reactant conversion or product selectivity makes it difficult to judge the potential of a catalytic transformation.

*T. Utne, J. D. Garber, R. E. Jones, US Patent 3083236, 1963.*

3. The catalytic hydrogenation of HMF to tetrahydrofuran-dimethanol (THF-dimethanol) using Raney nickel as described by Sutherland *et al.* is better described as a stoichiometric rather than a catalytic reaction.

*T. J. Connolly, J. L. Considine, Z. Ding, B. Forsatz, M. N. Jennings, M. F. MacEwan, K. M. McCoy, D. W. Place, A. Sharma, K. Sutherland, Org. Process Res. Dev. 2010, 14, 459.*

4. The reported 90% yield of THF-dimethanol using Raney nickel catalysts after 1 h reaction time is not in line with the proposed kinetic model.

*V. Schiavo, G. Descotes, J. Mentech, Bull. Soc. Chim. Fr. 1991, 128, 704.*

5. The addition of one drop of HCl, as described in the experimental procedure for a catalytic transfer hydrogenation, is not a very exact number and may lead to erroneous results.

*R. C. Mebane, A. J. Mansfield, Synth. Commun. 2005, 35, 3083.*

6. The oxidation of 1,6-hexanediol using 80% nitric acid as reported by Nikolaeva *et al.* does not comply with the green chemistry and technology principles as large amounts of nitrogen dioxide are produced.

*N. V. Svetlakov, V. G. Nikitin, E. A. Nikolaeva, Russ. J. Org. Chem. **2007**, 43, 773.*

7. *Dimana ada kemauan, di situ ada jalan* (Where there is a will, there is a way).

*Indonesian proverb.*